

# CONTENTS

<b>Contents</b>	<b>ix</b>
<b>Abbreviations</b>	<b>xv</b>
<b>Summary</b>	<b>xix</b>
<b>Zusammenfassung</b>	<b>xxiii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Resorcin[4]arene Cavitands . . . . .	3
1.2 Monitoring Conformational Properties . . . . .	4
1.3 Vase-Kite1-Kite2 Kinetics and Thermodynamics . . . . .	5
1.4 Control of Conformational and Binding Properties . . . . .	7
1.5 Thesis Goal . . . . .	8
<b>2 Quinone Moieties as Redox-Active Wall Components</b>	<b>9</b>
2.1 Introduction . . . . .	10

2.2	Tetraquinone Cavitannds . . . . .	11
2.2.1	Synthesis . . . . .	11
2.2.2	Electrochemistry . . . . .	13
2.2.3	Redox Interconversion . . . . .	15
2.2.4	NMR Analysis . . . . .	15
2.2.5	Kite1-Kite2 Activation Parameters . . . . .	21
2.2.6	X-Ray Analysis . . . . .	31
2.2.7	DFT Calculations . . . . .	32
2.2.8	Binding Properties . . . . .	37
2.2.9	Summary and Conclusion . . . . .	38
2.3	Diquinone Cavitannds . . . . .	38
2.3.1	Synthesis . . . . .	39
2.3.2	Electrochemistry . . . . .	40
2.3.3	Redox Interconversion . . . . .	41
2.3.4	NMR Analysis . . . . .	42
2.3.5	Kite1-Kite2 Activation Parameters and Kite-Vase Thermodynamic Quantities . . . . .	47
2.3.6	X-Ray Analysis . . . . .	52
2.3.7	DFT Calculations . . . . .	54
2.3.8	Binding Properties . . . . .	57
2.3.9	Summary and Conclusion . . . . .	58
<b>3</b>	<b>Achieving Kite-Vase Switching Upon Reduction</b>	<b>59</b>
3.1	Introduction . . . . .	60
3.2	Towards <i>N</i> -Acetyl-Based Cavitannd . . . . .	61
3.3	Ester-Based Cavitannd . . . . .	63
3.3.1	Synthesis . . . . .	63
3.3.2	Redox Interconversion . . . . .	64
3.3.3	NMR Analysis . . . . .	64
3.3.4	DFT Calculations . . . . .	65
3.3.5	Summary and Conclusion . . . . .	66

---

3.4	<i>S/O</i> -Pyridine-Based Cavitands . . . . .	66
3.4.1	Synthesis . . . . .	67
3.4.2	Electrochemistry . . . . .	69
3.4.3	Redox Interconversion . . . . .	70
3.4.4	NMR Analysis . . . . .	71
3.4.5	DFT Calculations . . . . .	72
3.4.6	Summary and Conclusion . . . . .	73
3.5	Towards <i>N</i> -Pyridine-Based Cavitand . . . . .	73
3.5.1	Synthetic Efforts . . . . .	74
3.5.2	X-Ray Analysis . . . . .	75
3.5.3	Summary and Conclusion . . . . .	76
3.6	Pyridine-Based Cavitand . . . . .	76
3.6.1	Synthesis . . . . .	77
3.6.2	Electrochemistry . . . . .	78
3.6.3	Redox Interconversion . . . . .	78
3.6.4	NMR Analysis . . . . .	79
3.6.5	X-Ray Analysis . . . . .	81
3.6.6	DFT Calculations . . . . .	82
3.6.7	Summary and Conclusion . . . . .	85
3.7	Imidazole-Based Cavitand . . . . .	86
3.7.1	Synthesis . . . . .	86
3.7.2	Redox Interconversion . . . . .	88
3.7.3	NMR Analysis . . . . .	88
3.7.4	DFT Calculations . . . . .	90
3.7.5	Summary and Conclusion . . . . .	92
3.8	Carboxamide-Based Cavitand . . . . .	93
3.8.1	Synthesis . . . . .	93
3.8.2	Electrochemistry . . . . .	95
3.8.3	Redox Interconversion . . . . .	96
3.8.4	NMR Analysis . . . . .	97

3.8.5	X-Ray Analysis . . . . .	101
3.8.6	IR Analysis . . . . .	103
3.8.7	Binding Studies . . . . .	103
3.8.8	Binding Kinetics . . . . .	106
3.8.9	DFT Calculations . . . . .	110
3.8.10	Summary and Conclusion . . . . .	112
<b>4</b>	<b>Reducing Guest-Exchange Rates via Full Encapsulation</b>	<b>113</b>
4.1	Introduction . . . . .	114
4.2	Triptycenequinone-based Cavitand . . . . .	115
4.2.1	Synthesis . . . . .	115
4.2.2	Electrochemistry . . . . .	116
4.2.3	Redox Interconversion . . . . .	117
4.2.4	NMR Analysis . . . . .	118
4.2.5	X-Ray Analysis . . . . .	120
4.2.6	Binding Studies . . . . .	122
4.2.7	Summary and Conclusion . . . . .	125
4.3	Triptycenequinone-amide-based Cavitand . . . . .	125
4.3.1	Synthesis . . . . .	126
4.3.2	Redox Interconversion . . . . .	127
4.3.3	NMR Analysis . . . . .	128
4.3.4	X-Ray Analysis . . . . .	129
4.3.5	IR Analysis . . . . .	130
4.3.6	Binding Studies . . . . .	131
4.3.7	Guest Influence on Cavitand Stability Against Oxidation . .	132
4.3.8	Binding Kinetics . . . . .	136
4.3.9	Summary and Conclusion . . . . .	139
<b>5</b>	<b>Towards Functionalized Redox-Switchable Cavitands</b>	<b>141</b>
5.1	Introduction . . . . .	142
5.2	Retrosynthetic Analysis . . . . .	143

5.3	Synthetic Efforts . . . . .	145
5.4	DFT Study on Wall Precursor Reactivities . . . . .	150
5.5	Summary and Outlook . . . . .	152
<b>6</b>	<b>Investigation of Cavitand Dynamics</b>	<b>155</b>
6.1	Introduction . . . . .	156
6.2	Synthesis . . . . .	160
6.3	NMR Analysis . . . . .	162
6.4	X-Ray Analysis . . . . .	163
6.5	UV-Vis and Steady-State Fluorescence Data . . . . .	165
6.6	Time-Resolved Fluorescence Data . . . . .	166
6.6.1	Natural Donor and Acceptor Fluorescence Lifetimes . . . . .	167
6.6.2	Fluorescence Decays . . . . .	168
6.6.3	Fluorescence Anisotropy Decays . . . . .	170
6.7	Quantification of Donor-Only Ratios . . . . .	172
6.8	MD Simulations . . . . .	174
6.8.1	Linker Study . . . . .	175
6.8.2	Cavitand Study . . . . .	178
6.8.3	Fluorescence Decays from MD Data . . . . .	181
6.8.4	Fluorescence Anisotropy Decays from MD Data . . . . .	185
6.8.5	Distance Autocorrelation from MD Data . . . . .	186
6.9	Summary . . . . .	187
<b>7</b>	<b>Summary and Outlook</b>	<b>189</b>
<b>8</b>	<b>Experimental Part</b>	<b>197</b>
8.1	Synthesis Section . . . . .	198
8.2	Electrochemistry . . . . .	262
8.3	Binding Studies . . . . .	262
8.3.1	Titration Procedure . . . . .	262
8.3.2	Data Treatment . . . . .	263

8.4	MD Simulations . . . . .	264
8.4.1	Preparation of GAFF Input Files . . . . .	264
8.4.2	Preparation of CGenFF Input Files . . . . .	264
8.4.3	MD Run Parameters . . . . .	264
8.5	Time-Resolved Fluorescence Spectroscopy . . . . .	265
<b>A</b>	<b>Appendix</b> . . . . .	<b>267</b>
A.1	NMR Spectra of New Compounds . . . . .	268
A.2	X-Ray Analysis . . . . .	327
A.2.1	Tetraquinone Cavitand From (CH <sub>3</sub> ) <sub>2</sub> CO . . . . .	327
A.2.2	Pyridine-Based Cavitand from MeCN/CH <sub>2</sub> Cl <sub>2</sub> . . . . .	328
A.2.3	Triptycene-Based Cavitand from DMF . . . . .	332
A.2.4	Triptycene-Based Cavitand from Mesitylene- <i>d</i> <sub>12</sub> . . . . .	334
A.2.5	Diquinone Cavitand from DMF . . . . .	337
A.2.6	Diquinone Cavitand from Toluene . . . . .	340
A.2.7	Reduced Amide-Based Cavitand from Mesitylene- <i>d</i> <sub>12</sub> / Cy- clooctane . . . . .	344
A.2.8	Reduced Triptycenequinone-amide-based Cavitand from Me- sitylene- <i>d</i> <sub>12</sub> / CDCl <sub>3</sub> . . . . .	347
A.2.9	Quinone-Based Diiodocavitand from CDCl <sub>3</sub> . . . . .	349
A.2.10	<i>N</i> -Pyridine Cavitand Wall Precursor from CHCl <sub>3</sub> /EtOH . . . . .	352
	<b>References</b> . . . . .	<b>357</b>
	<b>Curriculum Vitae</b> . . . . .	<b>378</b>