

Contents

Acknowledgements	iii
Abstract (English/Deutsch)	v
List of Figures	xiv
List of Tables	xv
1 Introduction	1
1.1 What Is Effective Connectivity?	1
1.2 A Framework for Extracting Effective Connectivity of Neural Circuits	3
1.2.1 Activity measurements are required on the level of spike trains.	5
1.2.2 The generalized integrate-and-fire model can be used to define effective couplings between neurons.	6
1.2.3 Effective connectivity can be analyzed using measures from graph theory.	10
1.2.4 Effective connectivity is a useful measure in applications.	11
1.3 Previous Results	12
1.3.1 Known Physiological Connectivity of Cortical Circuits	12
1.3.2 Known Physiological Connectivity for Certain Invertebrate Nervous Systems	14
1.3.3 Applications of Effective Connectivity Measures on Multi-Neuron Recordings	15
1.4 Thesis Contributions	16
2 Gerhard, Pipa, Lima, Neuenschwander, Gerstner: Extraction of network topology from multi-electrode recordings: Is there a small-world effect?	19
2.1 Abstract	20
2.2 Introduction	20
2.3 Material and Methods	23
2.3.1 Experimental setup	23
2.3.2 Generalized Linear Models	24
2.3.3 Inferring effective connectivity	28
2.3.4 Graph-theoretical analysis	29
2.4 Results	30
2.4.1 Basic results	30

2.4.2	Cortical networks do not show scale-free structure.	32
2.4.3	Cortical networks show a small, but significant small-world structure.	33
2.4.4	Distance-dependent connectivity accounts for observed small-world structure.	33
2.4.5	Localized sub-sampling overestimates true small-world-ness.	35
2.5	Discussion	38
2.5.1	Effective connectivity using Generalized Linear Models	38
2.5.2	Biased estimation of small-world-ness for experiments with multiple electrodes	39
2.5.3	Future directions	41
2.6	Contributions	41
3	Gerhard, Kispersky, Gutierrez, Marder, Kramer, Eden: Successful reconstruction of a physiological circuit with known connectivity from spiking activity alone	43
3.1	Abstract	44
3.2	Author Summary	44
3.3	Introduction	45
3.4	Results	46
3.4.1	A point process model and a direct measure of the coupling filter can correctly infer the known STG connectivity.	46
3.4.2	Inferred connectivity with a point process model is robust to parameter changes and can be replicated across independent data sets.	51
3.4.3	An alternative definition of coupling strength based on Granger causality fails to reconstruct the known physiological connectivity.	53
3.4.4	A linear rate model is insufficient to reconstruct the circuit diagram.	53
3.4.5	The point process model predicts changing effective networks due to pharmacological manipulation.	57
3.5	Discussion	61
3.5.1	The crab stomatogastric nervous system as a model system for network inference	61
3.5.2	Limitations of previous approaches	62
3.5.3	The point process model as an example of a statistical inference procedure	64
3.5.4	Relevance to identification of large-scale networks	66
3.6	Materials and Methods	67
3.6.1	Experimental details	67
3.6.2	Point process model	68
3.6.3	Linear rate model	71
3.6.4	Definition of coupling strength	72
3.6.5	Granger causality analysis	73
3.7	Supplementary Information: Goodness-of-fit of multivariate point process models	74
3.8	Supplementary Information: Goodness-of-fit of the linear rate models	76
3.9	Supplementary Information: Variability of coupling strengths	77

3.10 Contributions	77
4 Lütcke, Gerhard, Zenke, Gerstner, Helmchen: Inference of neuronal network spike dynamics and topology from calcium imaging data	79
4.1 Abstract	79
4.2 Introduction	80
4.3 Results	82
4.3.1 A framework for simulations of neural network calcium imaging data . .	82
4.3.2 Analysis of spike inference from simulated calcium signals in individual neurons	86
4.3.3 Influence of indicator properties and parameter choices	88
4.3.4 Spike inference under conditions of indicator saturation	90
4.3.5 Temporal precision of spike inference	92
4.3.6 Simulation of large-scale network dynamics of spiking neurons	94
4.3.7 Inference of structural connectivity from network activity: Establishing an upper limit	95
4.3.8 Connectivity inference after imperfect spike train reconstruction	98
4.3.9 Identification of graph topology	98
4.4 Discussion	102
4.4.1 Potential and limits of spike inference	102
4.4.2 Inferring network structure from calcium imaging data	104
4.4.3 Future directions	106
4.5 Materials and methods	107
4.5.1 Simulation of single-neuron spike trains, calcium dynamics, and indicator fluorescence signals	107
4.5.2 Reconstruction of spike trains from calcium indicator signals	111
4.5.3 Comparison of original and reconstructed spike trains	112
4.5.4 Large-scale network simulation and detailed neuron model	112
4.5.5 Connectivity reconstruction based on coupled point process models . .	114
4.5.6 Cross-correlation analysis	115
4.5.7 Surrogate model of spike train reconstruction	115
4.5.8 Identification of graph topology	116
4.6 Contributions	116
5 Additional Results	117
5.1 Sensitivity Analysis for Estimation of Small-World-Ness From Effective Connectivity	117
5.2 Sensitivity Analysis for Estimation of Network Motifs From Effective Connectivity	120
5.3 Sub-Sampling Bias Of Small-World-Measures	123
5.4 Contributions	125
6 Discussion	127

A	Gerhard, Gerstner: Rescaling, thinning or complementing? On goodness-of-fit procedures for point process models and GLMs	131
A.1	Abstract	131
A.2	Introduction	132
A.3	Methods	133
A.3.1	Representations of neural activity	133
A.3.2	Goodness-of-fit tests for point processes	134
A.3.3	Creating surrogate point processes from time series	136
A.4	Results	137
A.4.1	Example: Inhomogeneous Poisson process	138
A.4.2	Example: Renewal process	139
A.4.3	Example: Inhomogeneous Spike Response Model	140
A.5	Discussion	141
A.6	Contributions	142
B	Gerhard, Szegletes: Spline- and Wavelet-based Models of Neural Activity in Response to Natural Visual Stimulation	143
B.1	Abstract	143
B.2	Introduction	144
B.3	Methods	145
B.3.1	Toy data	145
B.3.2	Real data set	148
B.4	Results	149
B.4.1	Toy data	149
B.4.2	Real data set	150
B.5	Discussion	151
B.6	Contributions	152
	Bibliography	175