

Contents

1	Introduction	1
1.1	Introduction	1
1.2	Summary	5
2	Detecting Invariant Manifolds, Attractors, and Generalized KAM Tori in Aperiodically Forced Mechanical Systems	9
2.1	Set-up	9
2.1.1	Geodesic transport barriers in phase space	11
2.1.2	Computation of invariant sets as transport barriers	14
2.2	Results	17
2.2.1	Proof of concept: Periodically forced Duffing oscillator	17
2.2.2	The aperiodically forced Duffing oscillator	23
2.3	Summary and conclusions	27
3	Geodesic Transport Barriers in Jupiter's Atmosphere: A Video-Based Analysis	30
3.1	Introduction	30
3.2	Set-up	32
3.3	Geodesic LCS theory	33
3.3.1	Parabolic LCSs	33
3.3.2	Elliptic LCSs	37
3.4	Unsteady transport barriers in the atmosphere of Jupiter	39
3.4.1	Prior work and present objectives	39
3.4.2	Video footage	39
3.4.3	Optical velocity field reconstruction	40
3.4.4	Validation of the reconstructed velocity field	45
3.4.5	Lagrangian advection	46

3.4.6	Unsteady zonal jet cores as parabolic LCSs	46
3.4.7	The Great Red Spot as a generalized KAM region	48
3.5	Summary and conclusions	50
4	A Spectral Clustering Approach to Lagrangian Vortex Detection	52
4.1	Introduction	52
4.2	Method	55
4.2.1	Input: A trajectory data set	55
4.2.2	Similarity graph construction	55
4.2.3	Graph sparsification	57
4.2.4	Spectral clustering	57
4.2.5	Estimating the number of clusters by eigenspace analysis	60
4.2.6	Retrieving clusters from matrix U by K-means clustering	60
4.2.7	Large-scale spectral clustering	61
4.3	Related previous work	63
4.3.1	The transfer-operator approach	63
4.3.2	Hierarchical partitioning of the transfer-operator	66
4.3.3	Application of Infomap to the transfer operator	66
4.3.4	Application of fuzzy clustering to a trajectory data set	67
4.4	Results	68
4.4.1	The periodically forced pendulum	69
4.4.2	Quasiperiodic Bickley jet	72
4.4.3	An ocean surface data set	73
4.4.4	The ABC flow	77
4.5	Summary and conclusions	80
4.A	Approximating Ncut	81
4.B	Bipartite spectral graph partitioning	82
5	Level Set Formulation of Two-dimensional Lagrangian Vortex Detection Methods	84
5.1	Background	84
5.1.1	Implicit boundary representation	84
5.1.2	Front evolution and level set theory	85
5.2	Variational level-set-based vortex detection	86
5.2.1	Stretching-based formulation	87
5.2.2	Graph-based formulation	88
5.3	Numerical results	90

5.3.1	Periodically forced pendulum	92
5.3.2	Jupiter's wind-velocity field	94
5.3.3	An ocean surface data set	96
5.4	Conclusion	98
5.A	Numerical aspects	98
5.B	Sparse Field Method	100
6	A Comparison of Lagrangian Methods for Coherent Structure Detection	102
6.1	Diagnostics for Lagrangian coherence	102
6.1.1	Finite-time Lyapunov exponent (FTLE)	103
6.1.2	Finite-Size Lyapunov Exponent (FSLE)	104
6.1.3	Mesochronic analysis	104
6.1.4	Trajectory length	105
6.1.5	Trajectory complexity	106
6.1.6	Shape coherence	106
6.2	Mathematical approaches to Lagrangian coherence	107
6.2.1	Geodesic theory of LCS	107
6.2.2	Rotationally coherent Lagrangian vortices	109
6.2.3	Coherent pairs of sets	112
6.2.4	Spectral clustering of trajectories	113
6.3	Method comparisons on three examples	115
6.3.1	Bickley jet	116
6.3.2	Two-dimensional turbulence	123
6.3.3	Wind field from Jupiter's atmosphere	129
6.4	Summary and conclusions	131
7	Bibliography	133