

Contents

Abstract	i
Kurzfassung	iii
1 Theoretical introduction	1
1.1 Strong-field physics	1
1.1.1 Keldysh parameter	1
1.1.2 The electric dipole approximation	3
1.1.3 Gauge transformations	4
1.1.3.1 Classical electromagnetism	4
1.1.3.2 Velocity gauge	4
1.1.3.3 Length gauge	5
1.1.4 Free-particle in an electromagnetic field: The Volkov solution	5
1.1.4.1 The Volkov solution in the velocity gauge	5
1.1.4.2 The Volkov solution in the length gauge	5
1.2 High-harmonic generation	6
1.2.1 The three-step model	7
1.2.2 High-harmonic spectroscopy	8
1.2.3 HHG in the strong-field approximation	9
1.2.4 The saddle-point method	13
1.2.4.1 Momentum-saddle-point approximation	13
1.2.4.2 Full saddle-point approximation	15
1.2.5 HHG driven by two-color circularly-polarized fields	17
1.2.5.1 Polarization as a parameter in the HHG	17
1.2.5.2 Selection rules	19
1.2.5.3 Saddle-point analysis of BHHG	20
2 Experimental details	25

2.1 Laser source	25
2.2 The HHG chamber	26
2.3 Sample delivery system for liquid systems	28
2.4 Sample-recovery system	28
2.5 High-harmonic generation driven by mid-IR pulses	29
2.6 Optical setups for BHHG and transient-grating BHHG	30
2.6.1 BHHG	30
2.6.1.1 Mach-Zehnder-type interferometric setup	30
2.6.1.2 In-line setup	31
2.6.1.3 Comparison of the two setups	33
2.6.2 HHG polarization measurement	33
2.6.3 Experimental setup for bi-circular transient grating spectroscopy (TG-BHHS)	33
3 BHHS as a probe of the symmetry of atomic orbitals	37
3.1 Introduction	37
3.2 Experimental details	40
3.3 Phase-matching considerations	41
3.4 Helicity asymmetry in neon and argon	43
3.5 Theoretical modelling of high-harmonic emission from <i>p</i> -orbitals in BCCP fields	46
3.5.1 TDSE calculations	46
3.5.2 Strong-field approximation with the saddle-point method	48
3.5.2.1 Ionization matrix element	49
3.5.2.2 Recombination matrix elements	50
3.5.2.3 Saddle-point analysis	52
3.5.3 Strong-field approximation with a "simplified" saddle-point method	52
3.6 Analysis	53
3.6.1 Helicity asymmetry in bi-circular high-harmonic generation	54
3.6.1.1 Propensity rules in the photorecombination step	54
3.6.1.2 Propensity rule in the ionization step	57
3.6.1.3 Combined results	57
3.6.2 Intensity dependence of the helicity asymmetry	60
3.7 Conclusion	61
4 Dynamical symmetry breaking in molecular systems	63

4.1	Experimental details	64
4.2	Bi-circular HHG in non-isotropic media	65
4.2.1	Experimental details	65
4.2.2	Theoretical analysis	67
4.2.2.1	Recombination matrix elements	68
4.2.2.2	Analysis of BHHG in an aligned ensemble	69
4.2.2.3	Discussion	72
4.3	Symmetry-breaking on an intra-molecular level	74
4.3.1	Experimental results	74
4.3.2	Remarks	75
5	Chiral discrimination through bi-elliptical high-harmonic spectroscopy	79
5.1	Introduction	79
5.2	Experimental details	82
5.2.1	Experimental setup	82
5.2.2	Gas delivery system	83
5.3	Experimental results	83
5.3.1	Chiral HHG in the "bi-circular-perpendicular" configuration	84
5.3.1.1	Data acquisition details and image analysis	84
5.3.1.2	Circular dichroism in the "bi-circular-perpendicular" arrangement	85
5.3.1.3	Laser parameters dependence	90
5.3.2	Chiral HHG in the "bi-circular-parallel" configuration	92
5.3.3	Chiral HHG in the "bi-elliptical" configuration	94
5.4	Theoretical description of CD in bi-circular fields	95
5.4.1	Mechanism 1: Lorentz-force mediated chiral asymmetry	96
5.4.2	Mechanism 2: Chiral dichroism induced by laser-driven dynamics in the cation	101
5.4.2.1	Evaluation of the ionization matrix elements using partial Fourier transform theory	103
5.4.2.2	Evaluation of the propagation matrix elements	106
5.4.2.3	Recombination amplitudes	107
5.4.2.4	Enantiomer-specific matrix elements. Grid discretization scheme	107
5.4.2.5	Quantum chemical calculations	111
5.5	Discussion	115

6	Time-resolved circular dichroism study of the ultrafast photodissociation of 2-iodobutane	119
6.1	Experimental design and realization	121
6.2	Photodissociation dynamics of 2-iodobutane in the \tilde{A} -band	127
6.2.1	Potential energy curves of the ground and the first two excited states of 2-iodobutane	127
6.2.2	Previous work on the photodissociation dynamics of 2-iodobutane	130
6.2.3	Wavepacket calculation of the \tilde{A} -band vibrational dynamics	131
6.3	Experimental results	132
6.3.1	Experimental details	132
6.3.2	Transient grating optimization	133
6.3.3	Characterization of the UV-induced photodissociation dynamics in 2-iodobutane	135
6.3.4	Time-resolved circular dichroism measurement in 2-iodobutane	140
6.3.4.1	Circular dichroism in the static regime	140
6.3.4.2	Time-dependent CD signal	141
6.3.5	Discussion	143
6.4	Large-amplitude internal rotation in the <i>sec</i> -butyl radical	145
7	Conclusion	153
	Curriculum Vitae	157
	Acknowledgements	160
	References	165