

Contents

| | |
|--|-----------|
| Zusammenfassung | I |
| Abstract | V |
| 1 Introduction | 1 |
| 1.1 Non-covalent Interactions | 2 |
| 1.1.1 Polar Interactions | 2 |
| 1.1.2 Hydrogen Bonds | 2 |
| 1.1.3 Non-Polar Interactions | 3 |
| 1.1.4 Hydrophobic Interactions | 3 |
| 1.1.5 Additivity, Solvent Effect and Enthalpy-Entropy Compensation | 4 |
| 1.2 Biomolecules in the Gas Phase: Relation to the Solution-Phase | 5 |
| 1.3 Considerations for the Analysis of Non-Covalent Complexes Using ESI-MS | 8 |
| 1.3.1 Sample Preparation & Solvent Compatibility | 8 |
| 1.3.2 Desolvation Conditions | 8 |
| 1.3.3 Electrochemistry in the ESI Capillary | 8 |
| 1.3.4 Ionization of the Analyte and Transfer to the Gas Phase | 9 |
| 1.3.5 Chemical Equilibrium & Quantification | 9 |
| 1.4 Characterization of Oligomeric Biomolecular Complexes Using Native MS | 11 |
| 1.5 Characterization of High Molecular Weight Capsids | 14 |
| 1.6 Motivation and Scope of this Thesis | 15 |
| 2 Overview of Methods for the Analysis of non-covalent Biomolecular Complexes | 19 |
| 2.1 Fundamentals - Mass Spectrometry in the Biosciences | 20 |
| 2.2 Principles & Practice of Electrospray Ionization | 20 |
| 2.3 Mechanisms for Molecular Ion Formation in ESI-MS | 22 |

| | | |
|----------|---|-----------|
| 2.4 | Nanoelectrospray Ionization | 25 |
| 2.5 | Analysis and Detection of Ions in a Mass Spectrometer | 26 |
| 2.6 | Principles of Mass Filtering in a Quadrupole | 28 |
| 2.7 | Hybrid Quadrupole Time-of-Flight (Q-ToF) Mass Spectrometer | 29 |
| 2.8 | Gas Phase Fragmentation | 30 |
| 2.9 | Ion Mobility Spectrometry - Mass Spectrometry (IMS-MS) | 32 |
| 2.10 | Matrix - Assisted Laser Desorption/Ionization Mass Spectrometry | 35 |
| 2.10.1 | The MALDI Process | 35 |
| 2.10.2 | Ionization in Biological MALDI-MS | 35 |
| 2.10.3 | High-Mass Detection in MALDI-MS | 36 |
| 2.11 | Combining Chemical Cross-linking with High-Mass MALDI-MS | 38 |
| 2.12 | Overview of Traditional Solution Phase Methods to Study Biomolecular Structure and Dynamics | 41 |
| 2.12.1 | X-ray Crystallography | 41 |
| 2.12.2 | Nuclear Magnetic Resonance (NMR) | 41 |
| 2.12.3 | Calorimetric Methods | 41 |
| 2.12.4 | Surface Plasmon Resonance (SPR) | 42 |
| 3 | Insight Into Signal Response of Protein Ions in Native ESI-MS from the Analysis of Model Mixtures of Covalently Linked Protein Oligomers | 43 |
| 3.1 | Introduction | 45 |
| 3.1.1 | Materials and Methods | 48 |
| 3.1.2 | Buffer Exchange | 48 |
| 3.1.3 | Surface Plasmon Resonance (SPR) Experiments | 49 |
| 3.1.4 | Mass Spectrometry | 50 |
| 3.1.5 | Data Processing | 51 |
| 3.2 | Results and Discussion | 51 |
| 3.2.1 | Individual MBP _n Proteins Exhibit Linear Signal Response in Native ESI-MS | 51 |
| 3.2.2 | Evidence of Ion Suppression from the Analysis of Equimolar Mixtures of MBP _n s | 54 |
| 3.2.3 | Mechanistic Suggestions for the Observed Ion Suppression in Equimolar Mixtures of MBP _n s | 56 |

| | | |
|----------|---|-----------|
| 3.2.4 | Changing the Ratio of the Individual Components in Binary Mixtures of MBP _n s Results in a Loss of the Proportionality between Signal Response and Concentration | 62 |
| 3.2.5 | Determination of a Non-covalent Dissociation Constant Using Native ESI-MS | 64 |
| 3.3 | Conclusions | 67 |
| 4 | High-Mass MALDI-MS Analysis for the Investigation of Protein Encapsulation within an Engineered Capsid Forming Protein | 69 |
| 4.1 | Introduction | 71 |
| 4.2 | Materials and Methods | 73 |
| 4.2.1 | Chemical Cross-linking | 74 |
| 4.2.2 | Mass Spectrometry | 74 |
| 4.3 | Results and Discussion | 75 |
| 4.3.1 | Determination of the Subunit Stoichiometry | 75 |
| 4.3.2 | Understanding the Encapsulation of GFP(+36) by AaLS-13 Capsids | 78 |
| 4.4 | Conclusion and Outlook | 83 |
| 5 | Native ESI-MS reveals heterohexamer formation of co-expressed CcmK isoforms and spontaneous assembly of shell subunits in β-carboxysomes | 85 |
| 5.1 | Introduction | 87 |
| 5.2 | Experimental Methods | 91 |
| 5.2.1 | Buffer Exchange | 91 |
| 5.2.2 | Determination of the Protein Concentration | 91 |
| 5.2.3 | Protein Denaturation | 91 |
| 5.2.4 | Chemical Cross-linking Protocol | 92 |
| 5.2.5 | Subunit Exchange Experiment | 92 |
| 5.2.6 | Mass Spectrometry | 92 |
| 5.2.7 | Ion Mobility Mass Spectrometry | 93 |
| 5.2.8 | MALDI Mass Spectrometry | 93 |
| 5.3 | Results and Discussion | 94 |
| 5.3.1 | Characterization of CcmK Isoforms in Solution Using ESI-MS | 94 |
| 5.3.2 | No Subunit Exchange in Mixtures of Individually Expressed CcmK _i Isoforms | 95 |

| | | |
|----------|---|------------|
| 5.3.3 | Co-expression of CcmK _i Isoforms Results in Heterohexamer Formation | 98 |
| 5.3.4 | Separately Expressed ¹⁵ N-CcmK1 and ¹³ C-CcmK1 do not Form Heterocomplexes | 101 |
| 5.3.5 | Studying the Spontaneous Assembly of CcmK Hexamer Subunits in β -Carboxysomes | 103 |
| 5.3.6 | Characterization of the Residual Shell Components Involved in the Formation of the Icosahedron: CcmP, CcmO and CcmL | 105 |
| 5.3.7 | Binary Mixtures of Shell Building Blocks are Insufficient for Capsid Fragment Formation | 108 |
| 5.3.8 | Co-expression of Shell Components Leads to Formation of Assembly Intermediates of Mixed Nature | 110 |
| 5.4 | Conclusion and Outlook | 111 |
| 6 | Aryl bis-sulfonamides bind to the active site of a homotrimeric isoprenoid biosynthesis enzyme IspF and extract the essential divalent metal cation cofactor | 113 |
| 6.1 | Introduction | 115 |
| 6.2 | Materials and Methods | 118 |
| 6.2.1 | Materials | 118 |
| 6.2.2 | Experimental | 119 |
| 6.3 | Results and Discussion | 125 |
| 6.3.1 | Analysis of <i>AtIspF</i> under Denaturing and Native-like Conditions Using Native ESI-MS | 125 |
| 6.3.2 | Binding of the Natural Ligands, CDP-MEP and Zn ²⁺ to <i>AtIspF</i> | 127 |
| 6.3.3 | Aryl Bis-sulfonamide Inhibitors Bind to <i>AtIspF</i> Trimer and Extract Zn ²⁺ Cation from the Active Site | 130 |
| 6.3.4 | Docking of Bis-sulfonamide Ligands | 136 |
| 6.3.5 | CDP-ME and Aryl Bis-sulfonamide Inhibitors Compete for the Same Binding Site in <i>AtIspF</i> | 139 |
| 6.3.6 | Suggested Mechanism of Bis-sulfonamide Inhibitors Binding to <i>AtIspF</i> | 141 |
| 6.4 | Conclusions | 142 |
| 7 | Summary and Outlook | 145 |
| 7.1 | Summary | 146 |

| | | |
|-------------------|--|------------|
| 7.2 | Outlook | 150 |
| References | | 153 |
| A | Supporting Information | 177 |
| A.1 | Insight Into Signal Response of Protein Ions in Native ESI-MS from the Analysis of Model Mixtures of Covalently Linked Protein Oligomers | 177 |
| A.1.1 | Derivation of Binary Mixtures Models | 178 |
| A.1.2 | Constant Total Concentration in a Binary Mixture of Components | 178 |
| A.1.3 | Calibration of the TWIMS Cell and Theoretical Estimation of the CCS of the Protein-N ₂ Pair | 180 |
| A.1.4 | Derivation of the Non-covalent Dissociation Constant Equilibrium Model for Anti MBP Antibody Binding MBP | 181 |
| A.2 | Native ESI-MS Reveals Heterohexamer Formation of Co-expressed CcmK Isoforms and Spontaneous Assembly of Shell Subunits in β -Carboxysomes | 184 |
| A.2.1 | Characterization of CcmK _i Isoforms | 184 |
| A.2.2 | Mixtures of Different Shell Bricks Involved in the Formation of the Carboxysome Shell | 189 |
| A.3 | Aryl Bis-sulfonamides Bind to the Active Site of a Homotrimeric Isoprenoid Biosynthesis Enzyme IspF and Extract the Essential Divalent Metal Cation Cofactor | 205 |
| A.3.1 | Molecular Weight of <i>AtIspF</i> Measured by ESI-MS | 205 |
| A.3.2 | Mixing Scheme of the Titration Experiments | 205 |
| B | Abbreviations | 213 |
| C | Contributions | 215 |
| D | Curriculum Vitae | 219 |