

# Contents

Abstract	3
Zusammenfassung	5
1 Introduction	9
2 Mathematical introduction	17
2.1 Hopf algebras	17
2.2 $q$ -deformed universal enveloping algebras	23
2.3 Quasi-triangular structure of $\mathcal{U}_h(\mathfrak{g})$	27
2.4 $\mathcal{U}_h(\mathfrak{sl}(3))$ as a quantum double	33
3 Maximally extended $\mathfrak{sl}(2 2)$ as a Quantum Double	37
3.1 Maximally extended $\mathcal{U}_h(\mathfrak{psl}(2 2))$	37
3.1.1 Overview of the algebras	37
3.1.2 Hopf structure of the centrally extended algebra	39
3.1.3 Hopf structure of the maximally extended algebra	41
3.1.4 Special features	43
3.2 Extending the algebra	45
3.2.1 Dual of the centrally extended Borel subalgebra	46
3.2.2 Extending the positive Borel subalgebra	49
3.2.3 Doubling the extended subalgebra	55
3.3 R-matrix	57
3.3.1 Basis	57
3.3.2 Computation	58
3.3.3 Chevalley–Serre form	60
3.4 Classical limit	62
3.5 Conclusions and discussion	64
4 Maximally extended $\mathfrak{sl}(2 2)$ as a contraction	65
4.1 Deformation of 3D Poincaré as a contraction	66
4.1.1 Hopf algebra	66
4.1.2 Universal R-matrix	69
4.2 Relation to kappa-Poincaré	71
4.2.1 Comparison	71
4.2.2 Classical limit	73
4.2.3 Higher-dimensional kappa-Poincaré algebras	74
4.2.4 R-matrix and 3D scattering problem	75
4.3 Maximally extended $\mathfrak{sl}(2 2)$ from $\mathfrak{d}(2, 1; \epsilon) \times \mathfrak{sl}(2)$	77

4.3.1	Lie superalgebra $\mathfrak{d}(2, 1; \epsilon)$ . . . . .	78
4.3.2	q-deformation of $\mathfrak{d}(2, 1; \epsilon)$ . . . . .	78
4.3.3	Contraction limit . . . . .	81
4.4	Conclusion . . . . .	83
<b>5</b>	<b>Maximally extended <math>\mathfrak{sl}(2 2)</math> and the deformed Hubbard model</b>	<b>85</b>
5.1	The deformed Hubbard model algebra . . . . .	85
5.2	Braiding of the maximally extended $\mathfrak{sl}(2 2)$ . . . . .	86
5.3	Quotient . . . . .	87
5.4	Conclusions . . . . .	93
<b>6</b>	<b>Conclusion</b>	<b>95</b>
	<b>Acknowledgements</b>	<b>99</b>
<b>A</b>	<b>Details of the R-matrix calculation</b>	<b>101</b>