

Contents

Acknowledgements	v
Abstract (English/French)	vii
List of figures	xii
List of tables	xv
1 Introduction	1
1.1 2D transition metal dichalcogenides	1
1.2 The Peierls instability	3
1.3 Outline	5
2 Methodology	7
2.1 Density functional theory	7
2.2 Density-functional perturbation theory	14
2.3 Dynamical mean-field theory	16
2.4 Quasiparticle and excitonic effects	19
3 Crystal field effects in two-dimensional transition metal dichalcogenides	23
3.1 Introduction	24
3.2 Methodology	26
3.2.1 Wannierization and crystal field parameters	26
3.2.2 Computational details	28
3.3 The case of TaS ₂	29
3.3.1 spd, pd, and d models	29
3.3.2 Semi-core and high-energy states	37
3.3.3 Spin-orbit coupling	38
3.4 Crystal field and the relative stability of the 1 <i>T</i> and 1 <i>H</i> phases	39
3.5 Trends across the periodic table	44
3.6 Relation to previous work	47
3.7 Conclusion	48
3.8 Appendix	49
3.8.1 Higher-energy bands of 1 <i>T</i> -TaS ₂	49
3.8.2 Band structures of the considered materials	50

4 A unified picture for lattice instabilities in metallic octahedral transition metal dichalcogenides	57
4.1 Appendix	65
4.1.1 Computational methodology	65
4.1.2 Wannier functions for $1T'$ -WS ₂	65
4.1.3 Wannier functions for $1T''$ -ReS ₂	66
4.1.4 Fermi surface for $1T$ -WS ₂	67
5 Charge density wave phase, Mottness, and ferromagnetism in two-dimensional $1T$-NbSe₂	69
5.1 Introduction	70
5.2 Computational methodology	71
5.3 Charge-density-wave phases	72
5.4 Mottness in the Star-of-David phase	74
5.5 Magnetic phases	77
5.6 Conclusions	79
5.7 Appendix	79
5.7.1 Assessment of the effect of spin-orbit coupling	79
5.7.2 Investigation of the 4×4 CCDW phase	80
5.7.3 Type II Wannier functions	80
6 Excitonic effects in two-dimensional TiSe₂ from hybrid density functional theory	83
6.1 Computational details	91
7 After-thoughts, conclusions, and outlook	93
Bibliography	112
Curriculum Vitae	113