

Table of contents

1. Introduction	1
1.1. The Endoplasmic reticulum (ER) as site of protein folding	1
1.1.1. Sub-compartments of the ER	1
1.1.2. Polypeptide entry into the ER	1
1.1.3. N-glycosylation of nascent polypeptides	3
1.1.4. Polypeptide binding to the ER lectin chaperones CNX and CRT	4
1.1.5. UGGT1 as ER quality control sensor	6
1.2. Protein export from the ER	6
1.3. ER-associated degradation (ERAD)	7
1.3.1. N-glycan processing and substrate recognition	7
1.3.2. Dislocation across the ER membrane	8
1.3.2.1. E3-ubiquitin ligases are the core of dislocation machineries	8
1.3.2.2. The dislocon	9
1.3.3. Polypeptide extraction and proteasomal degradation	10
1.4. ER stress and the unfolded protein response (UPR)	11
1.4.1. Proteostasis and UPR induction	11
1.4.2. Protein kinase RNA-like ER kinase (PERK)	12
1.4.3. Inositol-requiring protein 1 (IRE1)	13
1.4.4. Activating transcription factor 6 (ATF6)	14
1.4.5. Mechanisms of UPR induction	15
1.4.6. UPR in physiology and disease	16
1.5. Lysosomal degradation of the ER	17
1.5.1. Autophagy degrades cellular material within lysosomes	17
1.5.2. Activation of autophagy	18
1.5.2.1. Induction of autophagy by nutrient deprivation	18
1.5.2.2. Induction of autophagy by ER stress	19
1.5.3. Mechanisms of autophagy	19
1.5.3.1. Initiation and phagophore nucleation	19
1.5.3.2. Phagophore elongation	21
1.5.3.3. Phagophore closure	22
1.5.3.4. Autophagosome fusion with the lysosome and degradation of the cargo	22
1.5.4. Selective autophagy	23
1.5.4.1. Selective autophagy of the ER: from historical evidences to receptors discovery	24
1.5.4.2. ER-phagy receptors in mammals	25
1.5.5. Lysosomal degradation of misfolded proteins of the ER	26
1.6. Aim of the PhD thesis	29
1.6.1. UPR induction by misfolded proteins	29
1.6.2. Study of selective, receptor-mediated ER turnover	31
1.6.2.1. SEC62 regulates selective ER turnover during recovery from ER stress	31

1.6.2.2.	Mechanisms regulating clearance of proteasome-resistant protein polymers from the mammalian ER	32
2.	Results	35
2.1.	Chemical stresses fail to mimic the unfolded protein response resulting from luminal load with unfolded polypeptides	36
2.2.	Translocon component SEC62 acts in endoplasmic reticulum turnover during stress recovery	56
2.3.	ER-to-lysosome-associated degradation of proteasome-resistant ATZ polymers occurs via receptor-mediated vesicular transport	119
3.	Discussion	143
3.1.	UPR induction by misfolded proteins	144
3.1.1.	Chemical- vs misfolded proteins-induced UPR	144
3.1.2.	Mechanisms of UPR sensors activation by misfolded proteins	145
3.1.3.	Misfolded proteins' features and UPR induction	145
3.1.4.	UPR- and transcriptionally-independent responses to unfolded proteins	146
3.2.	RecovER-phagy	148
3.2.1.	Selective activation of ER turnover by CPA washout	148
3.2.2.	SEC62 role in protein translocation vs recovER-phagy	149
3.2.3.	Mechanisms of recovER-phagy	150
3.2.4.	SEC62 in cancer	152
3.3.	ERLAD of proteasome-resistant ATZ polymers	153
3.3.1.	Selective ER turnover induction on ER-load with ATZ polymers	153
3.3.2.	Dual degradative route for misfolded ATZ: proteasomal vs lysosomal degradation	153
3.3.3.	Requirements of ATZ delivery to endolysosome	155
3.3.4.	Selection of ATZ polymers within the ER lumen	156
3.3.5.	FAM134B in ER-phagy and ERLAD of ATZ polymers	157
3.4.	Mechanisms of ER delivery: recovER-phagy vs ERLAD of ATZ polymers vs ER-phagy	159
3.5.	Conclusions and future perspectives	160
4.	References	163
5.	Appendix	185
5.1.	<i>Curriculum vitae</i>	
5.2.	Published review articles	
6.	Acknowledgments	208